

**Continuous implicit method for the solution of general
second order ordinary differential equations**

¹A. O. Adesanya, ²T. A. Anake and ³G. J. Oghoyon
*Department of Mathematics,
Covenant University, Ota,
Ogun State, Nigeria.*

Abstract

Methods of collocation and interpolation were adopted to generate a continuous implicit scheme for the solution of second order ordinary differential equation. Newton polynomial approximation method was used to generate the unknown parameter in the corrector. This enables us to solve both initial and boundary value problems.

Keywords

Collocation, interpolation, continuous, implicit, Newton's polynomial, corrector.

1.0 Introduction

The second order ordinary differential equation of the form

$$y'' = f(x, y, y') \quad (1.1)$$

subject to $y(a) = \eta_0, y'(a) = \eta_1$ is called initial value problem. When the condition is of the form $y(a) = \eta_0, y(b) = \eta_2$ for $a \leq x \leq b$ it is called a boundary value problem, where f is a continuous function. Scientific and technological problems often lead to mathematical modeling of real life applications such as motion of projectiles or orbiting bodies, population growth, chemical kinetics and economic growth. Differential equation is often used to model the problems and most times these equations do not have analytic solution, hence an approximate numerical method is required to solve the problems. Equations (1.1) is conventionally solved by first reducing it to the system of first order ordinary differential equation and then one applies the various methods available for solving the system of the first order. This approach is extensively discussed in the literature and we cite few examples among others, [1], [2], [3], [4], [5], [6], [7]. Although this approach has tremendous success yet it has certain drawback. For instance, computer programs associated with the method are often complicated especially when the subroutines to supply the starting values for the methods result in longer computer time and more computational work. In addition Vigo – Aguiar and Ramos [8], stated that these methods do not utilize additional information associated with a specific ordinary differential equation, such as oscillatory nature of the solution.

¹Corresponding author:

¹e-mail address: torlar10@yahoo.com,

¹Telephone: +234-8086049694, +234-0803373199

Journal of the Nigerian Association of Mathematical Physics Volume 15 (November, 2009), 71 - 78

Continuous implicit method, A. O. Adesanya, T. A. Anake and G. J. Oghoyon *J of NAMP*

Block methods for numerical solution of the first order ordinary differential equations have been proposed by several researchers such as in [18, 20, 21 and 27]. Rosser [19] introduced the 3- point implicit block method based on integration formulae which is basically Newton's cote type. Zanariah et al [26], proposed 3 points implicit block method based on Newton's backward divided difference formula.

Considerable attention has been devoted to development of method to solve special second order ordinary differential equation of the type $y'' = f(x, y)$ directly without reducing it to system of first order. For instance [1],[9], [10], [11] and [12] among others. Hairer and Wanner [13] proposed Nystrom type method and stated other conditions for determining the parameter of the method. Chawla and Sharma [14] proposed a method due to Runge kutta method. Method of linear multistep method have been considered by Awoyemi and Kayode [15], and Kayode [23]. These methods are predictor corrector methods, although the implementation of the method in a pc mode yield good accuracy, the procedure is costly to implement. For instance, pc subroutine are very complicated to write, since they require special techniques for supplying the starting values and for varying the step size, which lead to longer time and more human effort. Jator and Li [16], proposed an order 5 method that was implemented without the need for either predictor or starting values from other methods. Jator [6] proposed an order 6 method based on the same method. Adesanya et al. [17] proposed a two step method for the general solution of second order which is self starting and adopt Newton's polynomial to generate the starting value. Awoyemi et al. [24] recently proposed a self starting Numerov method. This method solves both initial and boundary value solution of ordinary differential equation. Yahaya [12] constructed a Numerov method from a quadratic continuous polynomial solution. This process led to method applied to both initial and boundary value problems.

In this work, we propose a block method for three steps. This method adopt Newton's polynomial approximation to generate the starting value and solves both initial and boundary value problems.

2.0 Methodology

We first state the uniqueness theorem for higher order ordinary differential equation with initial value problem

Theorem 2.1

Let

$$u^n = f(x, u, u', \dots, u^{(n-1)}), u^k(x_0) = c_k \quad (2.1)$$

$k = 0, 1, \dots, n - 1$, where u are scalars

Let R be the region defined by the inequalities $x_0 \leq x \leq x_0 + a$, $|s_j - c_j| \leq b$, $j = 0, 1, \dots, n - 1$, ($a > 0, b > 0$). Suppose $f(x, s_0, \dots, s_{n-1})$ is defined in R and in addition

- (a) f is non negative and non decreasing in each x, s_0, \dots, s_{n-1} in R
- (b) $f(x, c_0, \dots, c_{n-1}) > 0$ for $x_0 \leq x \leq x_0 + a$
- (c) $c_k \geq 0, k = 1, 2, \dots, n - 1$

Then the initial value problem (1.1) has a unique solution in R . (see Wend [25]) for details). We consider an approximate solution to (1.1) in power series

$$y(x) = \sum_{j=0}^k a_j \phi_j(x) \quad (2.2)$$

$\phi_j = \phi^j, a_j, j = 0(1)2k - 1$ are constants to be determined. Consider a linear multistep method of the form

$$y(x) = \sum_{r=0}^{t-1} \phi_r(x) y_{n+r} + h^2 \sum_{r=0}^{m-1} \varphi_r(x) f_{n+r} \quad (2.3)$$

where $x = [x_n, x_{n+r}]$, $k =$ step length, $m =$ the distinct collocation point, t is the interpolation point, for our method, the step length $k = 3$

$$\phi_r(x) = \sum_{i=0}^{i+m-1} \phi_{i+1,r} p_i(x), r = 0, 1, 2, \dots, m-1 \quad (2.4)$$

$$h^2 \varphi(x) = \sum_{i=0}^{i+m-1} h^2 \varphi_{i+1,r}(x) p_i(x), r=0, 1, 2, \dots, m-1 \quad (2.5)$$

$$y(x_{n+r}) = y_{n+r}, r \in [0, 1, 2, \dots, t-1] \quad y''(x) = f_{n+r}, r = 0, 1, m-1 \quad (2.6)$$

To get $\phi_j(x)$ and $\varphi_j(x)$, According to Yahaya [12], Onumanyi arrived at matrix of the form $DC = I$, where I is an identity matrix of dimension $(t+m) \times (t+m)$.

$$D = \begin{pmatrix} 1 & x_n & x_n^2 & \cdot & & x_n^{t+i-1} \\ 1 & x_{n+1} & x_{n+1}^2 & \cdot & & x_{n+1}^{t+i-1} \\ \cdot & \cdot & \cdot & \cdot & & \cdot \\ 1 & x_{n+t-1} & x_{n+t-1}^2 & \cdot & & x_{n+t-1}^{t+m-1} \\ 0 & 0 & 2 & \cdot & & \cdot \\ \cdot & \cdot & \cdot & \cdot & & \cdot \\ \cdot & \cdot & \cdot & \cdot & & \cdot \\ 0 & 0 & 2 & \cdot & (t+m-1)(t+m-2)x_{m-2}^{t+m-2} & \cdot \end{pmatrix} \quad (2.7)$$

$$c = \begin{pmatrix} \alpha_{1,0} & \alpha_{1,1} & \cdot & \alpha_{1,t-1} & h^2 \varphi_{1,0} & \cdot & h^2 \varphi_{1,m-1} \\ \alpha_{2,0} & \alpha_{2,1} & \cdot & \alpha_{2,t-1} & h^2 \varphi_{2,0} & \cdot & h^2 \varphi_{2,m-1} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \alpha_{t+m,0} & \alpha_{t+m,1} & \cdot & \alpha_{t+m,t-1} & h^2 \varphi_{t+m,0} & \cdot & h^2 \varphi_{t+m,m-1} \end{pmatrix} \quad (2.8)$$

3.0 Development value for the unknown

Theorem 3.1

Assuming that $f \in C^{n+1}[a, b]$ and $x_k \in [a, b]$ for $k=0, 1, n$ are distinct values, then $f(x) = y(x) + R_n(x)$, where $y(x)$ is a polynomial that can be used to approximate $f(x)$ For Newton's polynomial

$$y(x) = a_0 + a_1(x-x_0) + a_2(x-x_0)(x-x_1) + \dots + a_n(x-x_0)(x-x_1)\dots(x-x_{n-1}) \quad (3.1)$$

$f(x) \cong y(x)$, $R_n(x)$ is the remainder and has the form

$$R_n(x) = \frac{f^{n+1}}{(x_{n+1})!} (x-x_0)(x-x_1)\dots(x-x_{n-1})(x-x_n) \quad (3.2)$$

(See Awoyemi et al. [24]) for details.

4.0 Development of three steps method

In developing the method with step length $k=3$, we consider

$$D = \begin{bmatrix} 1 & x_{n+1} & x_{n+1}^2 & x_{n+1}^3 & x_{n+1}^4 & x_{n+1}^5 \\ 1 & x_{n+2} & x_{n+2}^2 & x_{n+2}^3 & x_{n+2}^4 & x_{n+2}^5 \\ 0 & 0 & 2 & 6x_n & 12x_n^2 & 20x_n^3 \\ 0 & 0 & 2 & 6x_{n+1} & 12x_{n+1}^2 & 20x_{n+1}^3 \\ 0 & 0 & 2 & 6x_{n+2} & 12x_{n+2}^2 & 20x_{n+2}^3 \\ 0 & 0 & 2 & 6x_{n+3} & 12x_{n+3}^2 & 20x_{n+3}^3 \end{bmatrix} \quad (4.1)$$

This gives

$$\begin{aligned} \alpha_1 &= -t, \quad \alpha_2 = t + 1 \\ \varphi_0(t) &= \frac{h^2}{360} (-3t^5 + 10t^3 - 7t) \\ \varphi_1 &= \frac{h^2}{120} (3t^5 + 5t^4 - 20t^3 + 22t) \\ \varphi_2 &= \frac{h^2}{120} (-3t^5 + 10t^4 + 10t^3 + 60t^2 + 43t) \\ \varphi_3 &= \frac{h^2}{360} (3t^5 + 15t^4 + 20t^3 - 8t) \\ \alpha_1' &= -1, \quad \alpha_2' = 1 \\ \varphi_0'(t) &= \frac{h^2}{360} (-15t^4 + 30t^2 - 7) \\ \varphi_1'(t) &= \frac{h^2}{120} (15t^4 + 20t^3 - 60t^2 + 22) \\ \varphi_2'(t) &= \frac{h^2}{120} (-15t^4 + 40t^3 + 30t^2 + 120t + 43) \\ \varphi_3'(t) &= \frac{h^2}{360} (15t^4 + 60t^3 + 60t^2 - 8) \end{aligned} \quad (4.2)$$

where $t = \frac{x - x_{n+2}}{h}$. Evaluating (4.2) at x_{n+3} i.e. when $t = 1$ and substituting the result in (2.3)

$$\text{gives} \quad 12y_{n+3} - 24y_{n+2} + 12y_{n+1} = h^2(f_{n+3} + 10f_{n+2} + f_{n+1}) \quad (4.4)$$

(4.4) has order $p=4$ and error constant $cp^{+2} = -\frac{1}{240}$

Evaluating (4.2) at x_n i.e. when $t = -2$ and substituting the result in (2.3) gives

$$12y_{n+2} - 24y_{n+1} + 12y_n = h^2(f_{n+2} + 10f_{n+1} + f_n) \quad (4.5)$$

(4.5) has order $p = 4$ and error constant $cp^{+2} = -\frac{1}{240}$.

Evaluating (4.3) at $t = -2$ and substituting the result in (2.3) gives

$$360hy'_n - 360y_{n+2} + 360y_{n+1} = h^2(-24f_{n+3} + 9f_{n+2} - 414f_{n+1} - 127f_n) \quad (4.6)$$

Evaluating (4.3) at $t = -1$ and substituting the result in (2.3) gives

$$360hy'_{n+1} - 360y_{n+2} + 360y_{n+1} = h^2(127f_{n+3} - 546f_{n+2} + 471f_{n+1} - 952f_n) \quad (4.7)$$

Evaluating (4.3) at $t = 0$ and substituting the result in (2.3) gives

$$360hy'_{n+2} - 360y_{n+2} + 360y_{n+1} = h^2(7f_{n+3} - 66f_{n+2} - 129f_{n+1} + 8f_n) \quad (4.8)$$

Evaluating (4.3) at $t = 1$, and substituting the result in (2.3) gives

$$360hy'_{n+3} - 360y_{n+2} + 360y_{n+1} = h^2(-8f_{n+3} + 129f_{n+2} + 66f_{n+1} - 7f_n) \quad (4.9)$$

Solving (4.3), (4.4) and (4.5) using matrix inversion method, we obtain the block

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} y_{n+1} \\ y_{n+2} \\ y_{n+3} \end{bmatrix} + \begin{bmatrix} 0 & 0 & -1 \\ 0 & 0 & -1 \\ 0 & 0 & -1 \end{bmatrix} \begin{bmatrix} y_{n-1} \\ y_{n-2} \\ y_n \end{bmatrix} = h^2 \begin{bmatrix} \frac{19}{60} & \frac{-13}{120} & \frac{1}{15} \\ \frac{22}{15} & \frac{2}{15} & \frac{2}{15} \\ \frac{27}{10} & \frac{27}{40} & \frac{17}{60} \end{bmatrix} \begin{bmatrix} f_{n+1} \\ f_{n+2} \\ f_{n+3} \end{bmatrix} + \frac{h^2}{360} \begin{bmatrix} 0 & 0 & \frac{97}{360} \\ 0 & 0 & \frac{28}{45} \\ 0 & 0 & \frac{39}{40} \end{bmatrix} \begin{bmatrix} f_{n-1} \\ f_{n-2} \\ f_n \end{bmatrix} \quad (4.10)$$

$$-h \begin{bmatrix} 0 & 0 & -1 \\ 0 & 0 & -2 \\ 0 & 0 & -3 \end{bmatrix} \begin{bmatrix} y'_{n-1} \\ y'_{n-2} \\ y'_n \end{bmatrix}$$

Hence, from (4.10)

$$y_{n+1} = y_n + \frac{h^2}{360}(114f_{n+1} - 39f_{n+2} + 24f_{n+3} + 97f_n) + hy'_n \quad (4.11)$$

$$y_{n+2} = y_n + \frac{h^2}{45}(66f_{n+1} + 6f_{n+2} + 6f_{n+3} + 28f_n) + 2hy'_n \quad (4.12)$$

$$y_{n+3} = y_n + \frac{h^2}{120}(324f_{n+1} + 81f_{n+2} + 34f_{n+3} + 117f_n) + 3hy'_n \quad (4.13)$$

4.1 Development of the unknown for $k = 3$

Evaluate the first derivative of (3.1), and neglect a_4 and higher values of a i.e. Newton's polynomial of order 4, we obtain

$$12hy'_n = -25y_n + 48y_{n+1} - 36y_{n+2} + 16y_{n+3} - 3y_{n+4} \quad (4.14)$$

$$12hy'_{n+1} = -3y_n - 10y_{n+1} + 18y_{n+2} - 6y_{n+3} + y_{n+4} \quad (4.15)$$

$$12hy'_{n+2} = y_n - 8y_{n+1} + 8y_{n+3} - y_{n+4} \quad (4.16)$$

$$12hy'_{n+3} = -y_n + 6y_{n+1} - 18y_{n+2} + 10y_{n+3} + 3y_{n+4} \quad (4.17)$$

Making y_{n+4} the subject in (4.14) and substituting into (4.15)-(4.16) and solving for y_{n+1}, y_{n+2} and y_{n+3} gives

$$y_{n+1} = y_n + \frac{h}{72}(57y'_{n+1} - 10y'_{n+2} + 3y'_{n+3} + 27y'_n) \quad (4.18)$$

$$y_{n+2} = y_n + \frac{h}{9}(127y'_{n+1} + 2y'_{n+2} + 3y'_n) \quad (4.19)$$

$$y_{n+3} = y_n + \frac{h}{8}(9y'_{n+1} + 6y'_{n+2} + 3y'_{n+3} + 3y'_{n+n}) \quad (4.20)$$

Make y_{n+1} the subject in (4.5) and then substitute in (4.6)-(4.8) to solve for f_{n+1} , f_{n+2} and f_{n+3} , give

$$f_{n+1} = -f_n + \frac{1}{h} \left(\frac{-3}{2} y'_{n+1} + \frac{3}{2} y'_{n+2} + \frac{1}{6} y'_{n+3} - \frac{1}{6} y'_n \right) \quad (4.21)$$

$$f_{n+2} = 3f_n + \frac{1}{h} \left(3y'_{n+1} - 6y'_{n+2} + \frac{7}{3} y'_{n+3} + \frac{2}{3} y'_n \right) \quad (4.22)$$

$$f_{n+3} = 25f_n + \frac{1}{h} \left(\frac{39}{2} y'_{n+1} - \frac{69}{2} y'_{n+2} + \frac{17}{2} y'_{n+3} + \frac{13}{2} y'_n \right) \quad (4.23)$$

Comparing (4.20)-(4.23) with the second derivative of (3.1) gives

$$y'_{n+1} = \frac{-486}{14417} y'_n - \frac{504}{14417} hf_n \quad (4.24)$$

$$y'_{n+2} = \frac{16974}{14417} y'_n - \frac{1620}{14417} hf_n \quad (4.25)$$

$$y'_{n+3} = \frac{63485}{14417} y'_n - \frac{45228}{14417} hf_n \quad (4.26)$$

5.0 Numerical example

We test the efficiency of our scheme on linear and non linear second order differential equation.

Problem 5.1:

$$y'' - x(y')^2 = 0$$

$$y(0) = 1, y'(0) = \frac{1}{2}, h = 0.1/40$$

$$\text{Exact solution } y(x) = 1 + \frac{1}{2} \ln \left(\frac{2+x}{2-x} \right)$$

Grid point	Expected result	Calculated result	Error
0.0025	1.00125000065104	1.00125000014116	5.09882D-10
0.0050	1.002500000520835	1.002500000032393	4.88542D-09
0.0075	1.00375001757828	1.00375001139519	6.18308D-09
0.0100	1.00500004166729	1.00500001779432	2.38729D-08
0.0125	1.00625008138212	1.00625002292868	5.84534D-08
0.0150	1.00750014062974	1.00750005100468	8.96250D-08
0.0175	1.00875022331755	1.00875020629768	1.60840D-07
0.0200	1.01000033335333	1.01000007231814	2.61035D-07
0.0225	1.01125047464542	1.01125041188283	3.55817D-07
0.0250	1.01250065110271	0.012500613537337	3.75653D-07

Problem 5.2

$$y'' = 2y^3$$

$$y(0) = 1, y'(0) = -1, h = 0.1/40$$

$$\text{Exact solution } y(x) = \frac{1}{x}$$

Grid point	Expected result	Calculated result	Error
0.0025	0.997506234413965	0.997506800724006	-4.560D-07
0.0050	0.995024875621891	0.995029339533017	1.941D-06
0.0075	0.992555831265509	0.992564223860174	-8.397D-06
0.0100	0.99009900990099	0.990070873966239	2.814D-05
0.0125	0.98765432098754	0.98759291299131	6.140D-05
0.0150	0.985221674876847	0.985127023087557	9.465D-05
0.0175	0.982800982800983	0.982633524787924	1.675D-04
0.0200	0.980392156862745	0.980155071394184	2.371D-04
0.0225	0.97799511002445	0.977688418640018	3.067D-04
0.0250	0.975609756097561	0.975194774137825	4.149D-04

Problem 5.3

$$y'' = y + e^{3x}$$

$$y(0) = \frac{-3}{32}, y'(0) = \frac{-5}{32}, h = 0.1/40$$

$$\text{Exact solution } y(x) = \frac{4x - 3}{32 \exp(-3x)}$$

Grid point	Expected result	Calculated result	Error
0.0025	-0.094140915761848	-0.094140939393182	2.34D-08
0.0050	-0.094532404142338	-0.094532599228254	1.95D-07
0.0075	-0.094924451608388	-0.094924817272551	3.65D-07
0.0100	-0.095317044390700	-0.095317760663383	7.16D-07
0.0125	-0.095710168480980	-0.095710743379670	5.74D-07
0.0150	-0.096103809629113	-0.096109967248178	6.16D-06
0.0175	-0.09649533403163	-0.096494619870395	7.14D-07
0.0200	-0.096892584872264	-0.096896306302397	3.72D-07
0.0225	-0.097289689232184	-0.097285656289237	4.03D-06
0.0250	-0.097683251173919	-0.097685517015441	2.26D-06

References

- [1] J. D. Lambert (1973), computational methods in ordinary differential equation, John Wiley, New York.
- [2] S. O. Fatunla (1991): Block method for second order ordinary differential equation. International journal of computer mathematics, volume 41, issue 1 & 2, pg 55-63.

- [3] Jennings (1987): Matrix computation for engineers and scientists, John Wiley and Sons, A Wiley Inter-science publication, New York.
- [4] L. Brujnanao, D. Trigiante (1998): Solving differential problems by multistep initial and boundary value methods. Gordon and Breach science publishers, Amsterdam, Pg 280-299
- [5] P. Onumanyi, S. N Jator, U. W. Serisena (1999): Continuous finite difference approximation for solving differential equation. International journal of Computer Mathematics, volume 72, No 1, pg 65-79
- [6] S. N. Jator (2001): Improvement in Adams – Moulton method for the first order initial value problems. Journal of the Tennessee Academy of science, volume 76, No 2, pg 57-60
- [7] D. O. Awoyemi (2001): A new Smith-order algorithm for general secondary order ordinary differential equation. International journal of computer mathematics volume 77, pg 117-124
- [8] J. Vigo – Aguiar, H. Ramos (2006): variable step size implementation of multistep method for $y'' = f(x, y, y')$. Journal of computational and applied mathematics volume 192 page 114-131.
- [9] S. O. Fatunla (1995): A class of block method for second order initial value problems. International journal of computer mathematics, volume 55 issue 1 & 2 page 119-133.
- [10] P. Henrici (1962): Discrete variable method for Odes, John Wiley, New York, USA.
- [11] V. A. Aladeselu (2007): Improved family of block method for special second order initial value problems Journal of Nigerian Association of mathematical physics volume 11, pg 153-158.
- [12] Y. A Yahaya (2007): A note on the construction of Numerov method through a quadratic continuous polynomial for the general second order ordinary differential equation. Journal of Nigerian Association of mathematical physics, volume II, page 253-260
- [13] E. Hairer, G. Wanner, (1976): A theory for Nystrom methods, Numerische mathematic, volume 25 page 283-400.
- [14] M. M. Chawla, S. R. Sharma (1985): Families of three stage third order Runge Kutta-Nystrom method for $y'' = f(x, y, y')$, journal of the Australian mathematical society, volume 26, page 375-386
- [15] D. O. Awoyemi, S. J Kayode (2005): A maximal order collocation method for direct solution of initial value problems of general second order ordinary differential equation. Proceedings of the conference organized by the National mathematical centre, Abuja, Nigeria
- [16] S. J. Jator, J. Li, (2006): A self starting linear multistep method for a direct solution of the general second order initial value problems (to appear)
- [17] A. O. Adesanya, Anake, T. A., Bishop S. A. (2009). Two steps block method for the solution of general second order initial value problems of ordinary differential equation (to appear).
- [18] W. E. Milne (1953): Numerical solution of differential equation, Wiley, New York.
- [19] J. B. Rosser (1967): A Runge Kutta for all season, SIAM, page 417-452.
- [20] P. B. Worland (1976): Parallel method for the numerical solution of ordinary differential equation, IEEE trans. Computer C-25, page 1045-1048.
- [21] Z. Omar (199): Developing parallel block method for solving higher orders ODES directly. Thesis, University putra Malaysia
- [22] L. F. Sham pine, H. Watts (1969): Block implicit one step method, international journal of Math-comp (23) page 731-740.
- [23] Kayode, S. J. (2005): Some continuous method for the solution initial value problems of general second order ordinary differential equation. PhD thesis, Federal university of Technology, Akure, Ondo State, Nigeria
- [24] D. O. Awoyemi, A.O. Adesanya, S. N. Ogunyebi (2009): Construction of self Starting Numerov method for the solution of initial value problems of general second order ordinary differential equation. Int. j. Num. Math, University of Ado Ekiti, Vol. 4 No. 2, pp. 269-278
- [25] V.V. Wend (1969): Existence and Uniqueness of solution of ordinary differential equation. Proceedings of the American Mathematical Society, Vol. 23, No. 1, pp 27-23
- [26] Zanariah Abdul, Mohamed Bin, Zurni Omar (2006): 3 point implicit block method for solving ordinary differential equation. Bulletin of the Malaysian mathematical science (2), 29(1), pp. 23-31